

# On the Löwner, Minus, and Star Partial Orderings of Nonnegative Definite Matrices and Their Squares

Jerzy K. Baksalary

*Department of Mathematics  
Tadeusz Kotarbiński Pedagogical University  
PL-65-069 Zielona Góra, Poland*

*and  
Department of Mathematical Sciences  
University of Tampere  
P.O. Box 607  
SF-33101 Tampere, Finland*

and

*Friedrich Pukelsheim  
Institut für Mathematik  
Universität Augsburg  
D-8900 Augsburg, Germany*

Submitted by Ingram Olkin

---

## ABSTRACT

We investigate how the ordering of two Hermitian nonnegative definite matrices  $\mathbf{A}$  and  $\mathbf{B}$  relates to the ordering of their squares  $\mathbf{A}^2$  and  $\mathbf{B}^2$ , in the sense of the Löwner partial ordering, the minus partial ordering, and the star partial ordering. The condition that  $\mathbf{A}$  and  $\mathbf{B}$  commute appears essential in these investigations. We also give some comments on possible extensions of our results by replacing the squares  $\mathbf{A}^2$  and  $\mathbf{B}^2$  with the  $k$ th powers  $\mathbf{A}^k$  and  $\mathbf{B}^k$ .

---

## 1. INTRODUCTION

For Hermitian nonnegative definite matrices  $\mathbf{A}$  and  $\mathbf{B}$ , the Löwner partial ordering  $\stackrel{\text{L}}{\leq}$ , the minus partial ordering  $\bar{\leq}$ , and the star partial ordering  $\stackrel{*}{\leq}$

are defined as follows:

$$\mathbf{A} \stackrel{L}{\leq} \mathbf{B} \Leftrightarrow \mathbf{B} - \mathbf{A} = \mathbf{K}\mathbf{K}^* \quad \text{for some matrix } \mathbf{K}, \quad (1)$$

$$\mathbf{A} \stackrel{-}{\leq} \mathbf{B} \Leftrightarrow \mathbf{A}^- \mathbf{A} = \mathbf{A}^- \mathbf{B} \quad \text{for some } \mathbf{A}^- \in \mathbf{A}\{1\}, \quad (2)$$

$$\mathbf{A} \stackrel{*}{\leq} \mathbf{B} \Leftrightarrow \mathbf{A}^2 = \mathbf{A}\mathbf{B}, \quad (3)$$

where  $\mathbf{K}^*$  in (1) is the conjugate transpose of  $\mathbf{K}$ , and  $\mathbf{A}\{1\}$  in (2) denotes the set of generalized inverses of  $\mathbf{A}$ , i.e.,  $\mathbf{A}\{1\} = \{\mathbf{X} : \mathbf{A}\mathbf{X}\mathbf{A} = \mathbf{A}\}$ . The ordering (1) dates back to Löwner (1934). The orderings (2) and (3) are restrictions to Hermitian matrices of the general definitions introduced by Hartwig (1980) and Drazin (1978), respectively. Hartwig (1980) showed in addition that the minus ordering is equivalent to *rank subtractivity*,

$$\mathbf{A} \stackrel{-}{\leq} \mathbf{B} \Leftrightarrow r(\mathbf{B} - \mathbf{A}) = r(\mathbf{B}) - r(\mathbf{A}). \quad (4)$$

In view of Marsaglia and Styan (1974, p. 188) and Cline and Funderlic (1979, p. 195), an alternative form of (4) for Hermitian  $\mathbf{A}$  and  $\mathbf{B}$  is

$$\mathbf{A} \stackrel{-}{\leq} \mathbf{B} \Leftrightarrow \mathcal{R}(\mathbf{A}) \subseteq \mathcal{R}(\mathbf{B}) \quad \text{and} \quad \mathbf{A}\mathbf{B}^+\mathbf{A} = \mathbf{A}, \quad (5)$$

where  $\mathcal{R}(\cdot)$  stands for the range and  $\mathbf{B}^+$  is the Moore-Penrose inverse of  $\mathbf{B}$ . It is known that for nonnegative definite matrices, the partial orderings (1), (2), and (3) follow the implications

$$\mathbf{A} \stackrel{*}{\leq} \mathbf{B} \Rightarrow \mathbf{A} \stackrel{-}{\leq} \mathbf{B} \Rightarrow \mathbf{A} \stackrel{L}{\leq} \mathbf{B}; \quad (6)$$

cd. Baksalary, Kala, and Kłaczyński (1983, p. 84) and Hartwig and Styan (1987, Theorem 2.1).

The purpose of this paper is to compare the relations  $\mathbf{A} \stackrel{L}{\leq} \mathbf{B}$ ,  $\mathbf{A} \stackrel{-}{\leq} \mathbf{B}$ , and  $\mathbf{A} \stackrel{*}{\leq} \mathbf{B}$  with the corresponding relations involving  $\mathbf{A}^2$  and  $\mathbf{B}^2$ . Notice that on the cone of Hermitian nonnegative definite matrices, the relations  $\stackrel{L}{\leq}$ ,  $\stackrel{-}{\leq}$ , and  $\stackrel{*}{\leq}$ , defined as

$$\mathbf{A} \stackrel{L}{\leq} \mathbf{B} \Leftrightarrow \mathbf{A}^2 \stackrel{L}{\leq} \mathbf{B}^2,$$

$$\mathbf{A} \stackrel{-}{\leq} \mathbf{B} \Leftrightarrow \mathbf{A}^2 \stackrel{-}{\leq} \mathbf{B}^2,$$

$$\mathbf{A} \stackrel{*}{\leq} \mathbf{B} \Leftrightarrow \mathbf{A}^2 \stackrel{*}{\leq} \mathbf{B}^2,$$

specify further partial orderings. This is due to the fact that if  $\mathbf{A}$  and  $\mathbf{B}$  are Hermitian nonnegative definite matrices, then the equality between  $\mathbf{A}^2$  and  $\mathbf{B}^2$  implies the equality between their unique square roots  $\mathbf{A}$  and  $\mathbf{B}$ . Śtepniaĳ (1987) studied the problem of how the ordering  $\stackrel{L}{\leq}$  behaves on *sets* of matrices; see also Horn (1988) for an example disproving Śtepniaĳ's conjecture. In contrast, we here concentrate on *pairs* of matrices  $\mathbf{A}$  and  $\mathbf{B}$ , without any further algebraic structure. Moreover, we consider the orderings  $\stackrel{-}{\leq}$  and  $\stackrel{*}{\leq}$  in addition to  $\stackrel{L}{\leq}$ . In the final section, we give some comments on possible extensions of our results by replacing the squares  $\mathbf{A}^2$  and  $\mathbf{B}^2$  with the  $k$ th powers  $\mathbf{A}^k$  and  $\mathbf{B}^k$ .

## 2. RESULTS

Let  $\mathbf{A}$  and  $\mathbf{B}$  be two Hermitian nonnegative definite matrices. For the Löwner ordering it is well known that

$$\mathbf{A}^2 \stackrel{L}{\leq} \mathbf{B}^2 \Rightarrow \mathbf{A} \stackrel{L}{\leq} \mathbf{B};$$

cf. Davis (1963, p. 199) and Marshall and Olĳin (1979, p. 464). The converse implication fails to hold in general, as can be seen by taking

$$\mathbf{A} = \begin{pmatrix} 1 & 2 \\ 2 & 4 \end{pmatrix} \quad \text{and} \quad \mathbf{B} = \begin{pmatrix} 3 & 0 \\ 0 & 6 \end{pmatrix}; \quad (7)$$

see also a more general counterexample in Marshall and Olĳin (1979, p. 465). For the minus partial ordering, neither of the relations  $\mathbf{A} \stackrel{-}{\leq} \mathbf{B}$  and  $\mathbf{A}^2 \stackrel{-}{\leq} \mathbf{B}^2$  implies the other. The matrices in (7) form an example that  $\mathbf{A} \stackrel{-}{\leq} \mathbf{B}$  does not imply  $\mathbf{A}^2 \stackrel{-}{\leq} \mathbf{B}^2$ , whereas

$$\mathbf{A} = \begin{pmatrix} 1 & 2 \\ 2 & 4 \end{pmatrix} \quad \text{and} \quad \mathbf{B} = \begin{pmatrix} \sqrt{15} & 0 \\ 0 & \sqrt{30} \end{pmatrix}$$

illustrate that  $\mathbf{A}^2 \stackrel{-}{\leq} \mathbf{B}^2$  does not entail  $\mathbf{A} \stackrel{-}{\leq} \mathbf{B}$ . However, when  $\mathbf{A}$  and  $\mathbf{B}$

commute, then they more nearly behave like real (or complex) numbers, and relationships between orderings of matrices and orderings of their squares become feasible. For the three partial orderings (1), (2), and (3), the results are as follows.

**THEOREM 1** (Löwner partial ordering). *For Hermitian nonnegative definite matrices  $\mathbf{A}$  and  $\mathbf{B}$  consider the following:*

- (a<sub>1</sub>)  $\mathbf{A} \stackrel{L}{\leq} \mathbf{B}$ ,
- (b<sub>1</sub>)  $\mathbf{A}^2 \stackrel{L}{\leq} \mathbf{B}^2$ ,
- (c)  $\mathbf{AB} = \mathbf{BA}$ .

Then (a<sub>1</sub>), (c)  $\Rightarrow$  (b<sub>1</sub>), and (b<sub>1</sub>)  $\Rightarrow$  (a<sub>1</sub>).

**THEOREM 2** (Minus partial ordering). *For Hermitian nonnegative definite matrices  $\mathbf{A}$  and  $\mathbf{B}$  consider the following:*

- (a<sub>2</sub>)  $\mathbf{A} \stackrel{-}{\leq} \mathbf{B}$ ,
- (b<sub>2</sub>)  $\mathbf{A}^2 \stackrel{-}{\leq} \mathbf{B}^2$ ,
- (c)  $\mathbf{AB} = \mathbf{BA}$ .

Then any two of these statements imply the third statement.

**THEOREM 3** (Star partial ordering). *For Hermitian nonnegative definite matrices  $\mathbf{A}$  and  $\mathbf{B}$  consider the following:*

- (a<sub>3</sub>)  $\mathbf{A} \stackrel{*}{\leq} \mathbf{B}$ ,
- (b<sub>3</sub>)  $\mathbf{A}^2 \stackrel{*}{\leq} \mathbf{B}^2$ ,
- (c)  $\mathbf{AB} = \mathbf{BA}$ .

Then (a<sub>3</sub>)  $\Leftrightarrow$  (b<sub>3</sub>)  $\Rightarrow$  (c).

We may summarize (6) and the results above by the following schematic diagram in which  $\Rightarrow$  denotes the usual implication and  $\rightarrow$  denotes the implication which is valid under the commutativity condition  $\mathbf{AB} = \mathbf{BA}$ :

$$\begin{array}{ccccc}
 \mathbf{A} \stackrel{*}{\leq} \mathbf{B} & \Rightarrow & \mathbf{A} \stackrel{-}{\leq} \mathbf{B} & \Rightarrow & \mathbf{A} \stackrel{L}{\leq} \mathbf{B} \\
 \uparrow \downarrow & & \downarrow \uparrow & & \uparrow \downarrow \\
 \mathbf{A}^2 \stackrel{*}{\leq} \mathbf{B}^2 & \Rightarrow & \mathbf{A}^2 \stackrel{-}{\leq} \mathbf{B}^2 & \Rightarrow & \mathbf{A}^2 \stackrel{L}{\leq} \mathbf{B}^2
 \end{array}$$

### 3. PROOFS

In the course of the proofs we will use the following lemma on commutativity of functions defined on the linear space  $\mathfrak{H}_n$  of all  $n \times n$  Hermitian

matrices. If  $A \in \mathbb{H}_n$  has the spectral decomposition  $A = U\Lambda U^*$ , where  $U$  is a unitary matrix and  $\Lambda$  is the diagonal matrix with the diagonal elements  $\lambda_1, \dots, \lambda_n$  equal to the eigenvalues of  $A$ , then a function  $f : \mathbb{R} \rightarrow \mathbb{R}$  gives rise to the function  $f_n : \mathbb{H}_n \rightarrow \mathbb{H}_n$  by way of  $f_n(A) = Uf_n(\Lambda)U^*$ , where  $f_n(\Lambda)$  is the diagonal matrix with the diagonal elements equal to  $f(\lambda_1), \dots, f(\lambda_n)$ .

LEMMA. Let  $A \in \mathbb{H}_n$  and  $B \in \mathbb{H}_n$ , let  $\lambda_1, \dots, \lambda_a$  and  $\mu_1, \dots, \mu_b$  be the distinct eigenvalues of  $A$  and  $B$ , and let the functions  $f : \{\lambda_1, \dots, \lambda_a\} \rightarrow \mathbb{R}$  and  $g : \{\mu_1, \dots, \mu_b\} \rightarrow \mathbb{R}$  be one-to-one. Then  $AB = BA$  if and only if  $f_n(A)g_n(B) = g_n(B)f_n(A)$ .

The proof of this lemma follows by using the fact that the commutativity is a necessary and sufficient condition for two Hermitian matrices to admit spectral decompositions with the same unitary matrix.

*Proof of Theorem 1.* For three alternative proofs that (b<sub>1</sub>) implies (a<sub>1</sub>) see Davis (1983, p. 199) and Marshall and Olkin (1979, p. 464).

If  $A$  and  $B$  commute, then

$$A = UD_AU^* \quad \text{and} \quad B = UD_BU^* \tag{8}$$

for some unitary matrix  $U$  and nonnegative diagonal matrices  $D_A$  and  $D_B$ . Consequently,

$$A \preceq B \iff D_A \preceq D_B \iff D_A^2 \preceq D_B^2 \iff A^2 \preceq B^2. \quad \blacksquare$$

*Proof of Theorem 2.* For the proof that (a<sub>2</sub>) and (b<sub>2</sub>) imply (c) first notice that, in view of (5), condition (b<sub>2</sub>) is equivalent to

$$\mathcal{R}(A) \subseteq \mathcal{R}(B) \quad \text{and} \quad A^2(B^2)^+A^2 = A^2. \tag{9}$$

The equality  $AB^+A = A$  in (5) entails

$$(A^2B^+ - A)(A^2B^+ - A)^* = A^2(B^+)^2A^2 - A^2,$$

and then the equality in (9) yields

$$A^2B^+ = A = B^+A^2, \tag{10}$$

since  $(B^+)^2 = (B^2)^+$ . Applying the Lemma with  $f$  and  $g$  defined as  $f(x) = x^2$  and  $g(x) = 1/x$  when  $x \neq 0$  and  $g(0) = 0$  to (10) leads to the commutativity of  $A$  and  $B$ .

The proof of the remaining two parts follows straightforwardly from the fact that, in view of (5) and (8),

$$\mathbf{A} \bar{\leq} \mathbf{B} \Leftrightarrow \mathcal{R}(\mathbf{D}_A) \subseteq \mathcal{R}(\mathbf{D}_B) \text{ and } \mathbf{D}_A \mathbf{D}_B^+ \mathbf{D}_A = \mathbf{D}_A,$$

and

$$\mathbf{A}^2 \bar{\leq} \mathbf{B}^2 \Leftrightarrow \mathcal{R}(\mathbf{D}_A) \subseteq \mathcal{R}(\mathbf{D}_B) \text{ and } \mathbf{D}_A^2 (\mathbf{D}_B^2)^+ \mathbf{D}_A^2 = \mathbf{D}_A^2.$$

This shows that if (c) holds, then conditions (a<sub>2</sub>) and (b<sub>2</sub>) are equivalent: each of them is satisfied if and only if every diagonal element of  $\mathbf{D}_A$  is equal either to zero or to the corresponding element of  $\mathbf{D}_B$ . ■

*Proof of Theorem 3.* The statement “(a<sub>3</sub>) ⇒ (c)” is a direct consequence of the definition (3), and the statement “(a<sub>3</sub>) ⇔ (b<sub>3</sub>)” follows from the part “(a<sub>2</sub>), (c) ⇔ (b<sub>2</sub>), (c)” of Theorem 2, noting that (a<sub>3</sub>) ⇔ (a<sub>2</sub>), (c) [cf. Hartwig and Styan (1986, Theorem 2)] and, similarly, (b<sub>3</sub>) ⇔ (b<sub>2</sub>), (c). ■

#### 4. COMMENTS ON POSSIBLE EXTENSIONS

Professor Ingram Olkin asked about possible extensions of Theorems 1, 2, and 3 by replacing the squares of  $\mathbf{A}$  and  $\mathbf{B}$  with the values  $\varphi(\mathbf{A})$  and  $\varphi(\mathbf{B})$  of some more general function  $\varphi : \mathbb{H}_n \rightarrow \mathbb{H}_n$ . We comment here on this question in the particular situation when  $\varphi$  is the  $k$ th power function, that is, when conditions (b<sub>1</sub>), (b<sub>2</sub>), and (b<sub>3</sub>) are replaced by (b<sub>1</sub><sup>\*</sup>)  $\mathbf{A}^k \stackrel{L}{\leq} \mathbf{B}^k$ , (b<sub>2</sub><sup>\*</sup>)  $\mathbf{A}^k \bar{\leq} \mathbf{B}^k$ , and (b<sub>3</sub><sup>\*</sup>)  $\mathbf{A}^k \stackrel{*}{\leq} \mathbf{B}^k$ , respectively.

Our first observation is that the modified version of Theorem 1 remains true for any  $k > 1$ . The part “(b<sub>1</sub><sup>\*</sup>) ⇒ (a<sub>1</sub>)” is due to Löwner (1934) [cf. Marshall and Olkin (1979, p. 464)], and the part “(a<sub>1</sub>), (c) ⇒ (b<sub>1</sub><sup>\*</sup>)” can be established by similar arguments to those in the proof of Theorem 1; see also a related result of Man (1970, Corollary 2).

The second observation is that, using the same argument of simultaneous diagonalization of  $\mathbf{A}$  and  $\mathbf{B}$ , the parts “(a<sub>2</sub>), (c) ⇒ (b<sub>2</sub>)” and “(b<sub>2</sub>), (c) ⇒ (a<sub>2</sub>)” can be extended to “(a<sub>2</sub>), (c) ⇒ (b<sub>2</sub><sup>\*</sup>)” and “(b<sub>2</sub><sup>\*</sup>), (c) ⇒ (a<sub>2</sub>).” We were unable, however, to prove or disprove that  $\mathbf{A} \bar{\leq} \mathbf{B}$  and  $\mathbf{A}^k \bar{\leq} \mathbf{B}^k$  together imply the commutativity of  $\mathbf{A}$  and  $\mathbf{B}$  when  $k \neq 2$ .

Our final observation is again positive, viz. that the statement in Theorem 3 can be generalized to the form (a<sub>3</sub>) ⇔ (b<sub>3</sub>) ⇒ (c).

*This research was begun while the first author was visiting the University of Dortmund and the University of Augsburg and completed while he was a*

Visiting Professor of the Academy of Finland. The grants from the Alfred Krupp von Bohlen and Halbuch-Stiftung, the Schwerpunktprogramm der Deutschen Forschungsgemeinschaft "Anwendungsbezogene Optimierung und Steuerung," and the Academy of Finland are gratefully acknowledged. Partial support was also provided by the Polish Academy of Sciences Grant No. CPBP 01.01.2/1.

The authors are grateful to Professor Ingram Olkin, an Advisory Editor of the journal, for his comments which led to shorter proofs in Section 3 and the discussion in Section 4.

## REFERENCES

- Baksalary, J. K., Kala, R., and Kłaczyński, K. 1983. The matrix inequality  $M \geq B^*MB$ , *Linear Algebra Appl.* 54:77–86.
- Cline, R. E. and Funderlic, R. E. 1979. The rank of a difference of matrices and associated generalized inverses, *Linear Algebra Appl.* 24:185–215.
- Davis, C. 1963. Notions generalizing convexity for functions defined on spaces of matrices, in *Proc. Symp. Pure Math.* VII, Amer. Math. Soc., Providence, pp. 187–201.
- Drazin, M. P. 1978. Natural structures on semigroups with involution, *Bull. Amer. Math. Soc.* 84:139–141.
- Hartwig, R. E. 1980. How to partially order regular elements? *Math. Japon.* 25:1–13.
- Hartwig, R. E. and Styan, G. P. H. 1986. On some characterizations of the "star" partial ordering for matrices and rank subtractivity, *Linear Algebra Appl.* 82:145–161.
- Hartwig, R. E. and Styan, G. P. H. 1987. Partially ordered idempotent matrices, in *Proceedings of the Second International Tampere Conference in Statistics* (T. Pukkila and S. Puntanen, Eds.), Dept. of Mathematical Sciences, Univ. of Tampere, Tampere, Finland, pp. 361–383.
- Horn, R. A. 1988. Review of "Two orderings on a convex cone of nonnegative definite matrices," by C. Stępnia, *Math. Rev.* 88i:15040.
- Löwner, K. 1934. Über monotone Matrixfunktionen, *Math. Z.* 38:177–216.
- Man, F. T. 1970. Some inequalities for positive definite matrices, *SIAM J. Appl. Math.* 19:679–681.
- Marsaglia, G. and Styan, G. P. H. 1974. Equalities and inequalities for ranks of matrices, *Linear and Multilinear Algebra* 2:269–292.
- Marshall, A. W. and Olkin, I. 1979. *Inequalities: Theory of Majorization and Its Applications*. Academic, New York.
- Stępnia, C. 1987. Two orderings on a convex cone of nonnegative definite matrices. *Linear Algebra Appl.* 94:263–272.